

ZAŠTITA ULJANE REPICE

VREME PRIMENE

RAZLOG TRETIRANJA

Setva, depozitorima
tretriranjem u redovima

Zemljišne štetočine

Septembar-Oktobar

Repičina lisna osa (*Athalia rosae*)

Buvači (*Phyllotreta spp*, *Psylliodes
chrysocephala*)

Od faze 2 lista do 15 cm visine,
a kada su korovi u fazi 3-5 listova

Širkolisni korovi

Uskolisni korovi

Povećanje kvaliteta i stimulacija oplodnje

U proleće
po pojavi štetočina

Repičin sjajnik (*Meligethes aeneus*)

Repičina pipa (*Ceutorhynchus napi*,
C.pallidactylus)

Preventivno ili po pojavi prvih simptoma
tokom cvetanja

Crna pegavost (*Alternaria brassicae*)

Bela trulež (*Sclerotinia sclerotiorum*)

14 dana pre žetve

Desikacija


IZBOR PREPARATA

FAZA

Saturn Ultra 5–10 kg/ha



a.m. deltametrin
0,3–0,5 l/ha



a.m. klopuralid 300
0,2–0,3 l/ha

Flupisor 1 l/ha ili
Totem 1,5 l/ha



Wuxal® Ascofol 2 l/ha ili
Wuxal® Boron plus 2 l/ha

a.m. deltametrin 0,5 l/ha

ili

Bifenicus C 0,1–0,125 l/ha



Postulat 0,8–1,0 l/ha



Brener 1 l/ha

