

# ZAŠTITA KUPUSA

## VREME PRIMENE

## RAZLOG TRETIRANJA

### Pre rasađivanja

Dezinfekcija zemljišta

Uskolisni i širokolisni korovi

Štetočine u zemljištu

### Zaštita rasada

Lisne štetočine

Folijarna ishrana

### 8-10 dana po rasađivanju

Plamenjača i crna pegavost

Kupusari, buvači i vaši

Folijarna ishrana

### Formiranje rozete

Kupusari, buvači, vaši i sovice

Višegodišnji uskolisni korovi

Folijarna ishrana

### Formiranje glavice

Plamenjača i crna pegavost

Lisne vaši

Kupusari i sovice

Folijarna ishrana

## IZBOR PREPARATA

## FAZE

**Basamid® Granulat** 30–50 g/m<sup>2</sup>

**Stomp® Aqua** 3 l/ha



**Karate® 0,4 GR** 15 kg/ha

**Šiman 200 SP** 4 g/2–4 l vode



**Wuxal® Top P** 20 ml/2–4 l vode



**Teatar plus** 1 l/ha

**Sever** 1,5–2 kg/ha +  
**ES plus** 1 l/ha



**Wuxal® Ascofol** 2 l/ha  
**Wuxal® Boron plus** 2 l/ha



**Šiman 200 SP** 0,4 l/ha +  
**Zakon** 0,14–0,2 l/ha

**Pantera® 40 EC** 1–2 l/ha



**Wuxal® Boron plus** 2 l/ha

**Teatar plus** 1 l/ha

**Sonar** 0,12–0,14 kg/ha

**Zakon** 125–175 ml/ha +  
**ES plus** 1 l/ha



**Wuxal® Boron plus** 2 l/ha