

# ZAŠTITA KRASTAVCA

VREME PRIMENE	RAZLOG TRETIRANJA
Pre setve ili sadnje	Dezinfekcija zemljišta
Zaštita rasada	Bakterioze
	Povećanje otpornosti biljaka na stres
Prilikom rasađivanja	Štetočine u zemljištu
1-2 nedelje po rasađivanju	Plamenjača
	Bakterioze
	Aminokiselina organskog porekla
3-4 nedelje po rasađivanju	Plamenjača
	Pepelnica
	Folijarna ishrana
Formiranje vreže	Plamenjača, crna pegavost i pepelnica
	Siva trulež
	Mikroelementi
Početak cvetanja	Plamenjača i crna pegavost
	Pepelnica
	Stimulacija oplodnje i povećanje otpornosti biljaka na stres
Tokom zrenja (između berbi)	Plamenjača i crna pegavost
	Siva trulež
	Vaši i bela leptirasta vaš
	Žuti paučinar
	Folijarna ishrana

## IZBOR PREPARATA

## FAZE

**Basamid® Granulat** 30–50 g/m<sup>2</sup>



**Everest** 0,4%

**Wuxal® Ascofol** 3 l/ha



**Karate® 0,4 GR** 15 kg/ha



**Ovation** 2,5 kg/ha

**Everest** 0,4%

**Wuxal® Amino** 0,4%  
**Wuxal® Ascofol** 2 l/ha



**Kventin** 0,5 l/ha

**Neimar** 0,25–0,5 kg/ha ili  
**Nimrod®** 0,8 l/ha

**Wuxal® K 40** 3 l/ha



**Teatar plus** 1,2 l/ha

**Signum®** 1,5 kg/ha

**Wuxal® Microplant** 2 l/ha



**Kventin** 0,5 l/ha

**Neimar** 0,25–0,5 kg/ha

**Wuxal® Ascofol** 3 l/ha



**Teatar** 1,2 l/ha

**Enter** 1,5 l/ha

**Sivanto® prime** 0,75 l/ha

**Kanemite® 15 SC** 1 l/ha

**Wuxal® Aminocal** 0,25%

